

Self Help Guide



RSACC ♀
Rape and Sexual Abuse Counselling Centre
Darlington and County Durham

Contents

Self Help Guide	1
Introduction	3
What kind of reactions do people have to trauma?	4
Why do people have these reactions?	6
Thoughts	9
- Your mind's reaction	9
- Some ways you can deal with your thoughts	11
Feelings	14
- Your emotional reactions	14
- Some ways of dealing with emotional reactions	16
Meditation and Relaxation	18
Grounding step-by-step	20
Body	23
- Your body's reactions	23
- Some ways to look after your body	24
Behaviour	26
- Your actions	26
It's never too late to talk	30
Key reactions: Flashbacks	32
Key reactions: Self-harm	35
Breaking it down	38
Further resources	40

Self Help Guide

This guide has been put together to help survivors of rape, childhood sexual abuse and/or sexual abuse to understand and process their own personal reactions to their experience. These experiences all come under the category of sexual violence. Sexual violence can affect every part of you. It effects how you think, how you feel, how you interact with others, your ability and methods of coping.

Throughout this guide we use the word survivor. You may not feel like you're a survivor but that's exactly what people who live with the psychological and emotional effects of sexual violence are. They are survivors!

Everyones experience of sexual violence is unique to them. Whether it happened in childhood or as an adult the effects it has on survivors shouldn't be underestimated. Sexual violence is something that happens to someone. It is not who they are, it does not define them.

Sexual violence is something that happens to someone. It is not who they are, it does not define them.



This self help guide is designed to help people who have experienced any form sexual violence as the reactions and questions that follow can be similar. It is normal to feel a wide range of emotions from confusion, anger, upset and anxiety, along with many others including shock and disbelief but it can get better. Being gentle and patient with yourself is a good way to begin your recovery. Recovery takes time and patience and it is different for all survivors.

Working out which parts of the guide are most helpful to you is a step towards understanding yourself and your reactions. Each section of this guide explains some common reactions, followed by self-help suggestions. It can be helpful to write your own ideas next to our suggestions or to highlight suggestions you particularly like. As you go through the guide, try to think about real situations in which some of the ideas might work. The healthy ways of coping are always the best.

Introduction

It is possible to recover from your experiences. It can take time. Being patient and kind to yourself is a good way to start your recovery.

Many reactions associated with sexual violence are natural safety mechanisms. Your body and mind are reacting to a dangerous situation, they are doing this in an attempt to protect you from further harm:


- Fear and anger are protective feelings
- Mistrust and forgetfulness are protective thoughts
- Isolation and forming extra identities are protective behaviour

Your body will continue to be on high alert in an attempt to protect you from further harm. There is no knowing how long this will continue.

**Remember:
What you are feeling is a normal reaction.**

Choosing to start working on the healing process takes courage. Opening this booklet is a really good place to start.

Try to be patient and kind to yourself. Read this guide at your own pace: take breaks from it and try not to overwhelm yourself.



Remind yourself...

- It was never your fault
- Whatever your reactions, they are normal
- Many people don't tell anybody for many years
- Most perpetrators of rape or abuse are known to their victims
- You are not alone and there is support available

What kind of reactions do people have to trauma?

A trauma is an event or experience that causes your body or mind to shock itself into protection mode. When a person has experienced a shocking, unexpected or traumatic incident they are likely to develop deep emotional and physical shock or stress.

Reactions to these events can be profound and affect every aspect of a person.



Below are some of the effects survivors of sexual violence can experience.

Here are some typical reactions that people can experience after a trauma:

Physical

- Tension in the muscles
- Tiredness & exhaustion
- Diarrhoea
- Little desire to do anything
- Hyperactivity
- Dry mouth
- Sweating
- Headaches
- Dizzy spells/funny turns
- Unsteady breathing
- Tightness in the chest
- Increased, rapid heartbeat
- Other physical pains

Emotional

- Feelings of hopelessness
- Guilt
- Depression
- Insecurity
- Irritability
- Anxiety
- Moodiness
- Panic attacks
- Nightmares
- Poor memory
- Loss in self-confidence or concentration

Behavioural

- Increased smoking and/or drinking
- Over working, or not turning up to work
- Personal neglect
- Cutting or other self-harm
- Nail biting
- Nonstop talking
- Changes in eating
- Impulsiveness

This guide will use the word *'survivors'*.

It may not feel like it and you might think you have difficulty coping, but you have survived and you are surviving now.



Why do people have these reactions?

The human brain has evolved over thousands of years to protect and make sense of our world. It is capable of rational thought and intuitive or instinctual thoughts and actions.

When you are exposed to danger our brain switches to survival or protection mode. Its ability to rationalise is impaired and in a lot of cases shuts down.



Our senses are constantly sending signals to a part of our brain called the amygdala.

The amygdala is the brain's own personal security guard. It is there to protect and serve. It is not capable of rational thought and problem solving. If threats are found by the amygdala, signals are sent to the body to release protective hormones such as adrenaline and the stress hormone cortisol. It's this release of hormones that causes the survival response commonly known as **'fight or flight'**. This response makes us want to either fight, run away, freeze or flop. All of these are natural responses to a threatening situation.

Our body chooses the reaction most likely to protect us with the least harm.

It doesn't think about how you feel after. During sexual violence running away or fighting off the perpetrator isn't the safest option as it could lead to more physical harm. The other options of freeze or flop expose the survivor to the least immediate danger. Although people imagine the natural response would be to back, the reality is very different.

This all happens in a split second. Immediately after the amygdala begins the slower process of sending the threat signal to the rational part of our brain (the cortex or hippocampus). It is not until a few seconds later that we manage to think rationally about the threat. As the instinct for immediate survival overrides longer-term rational thought, fear can overwhelm rationality during trauma.

If there is a safe outcome (survival), the brain learns to use that reaction again.

Sometimes, this response can be used repeatedly in less and less risky situations. This can lead to a heightened state of awareness of risk, or to a feeling of numbness.



**I cannot
talk about it**


**If only
I hadn't...**

**They should
have protected
me**

**I'm not
coping**

**I have
no control**

**Nobody
understands**



Thoughts

Your mind's reactions

It is normal for you to keep thinking over your experience. Although this is usually unpleasant it is simply your mind's way of trying to make sense of what's happened. Along with this, your thoughts and feelings may alter or change in intensity. Everyone's experience is different but there are some similarities in how sexual violence affects us as human beings.

Angry thoughts

- They should have protected me
- No man can be trusted

Self-loathing

- I'm dirty and disgusting
- Nobody will ever understand me
- Nobody will ever love me

Feeling different

- Nobody understands
- Things will never be the same

>>



Flashback


A flashback is a memory suddenly, unexpectedly and vividly revisited. You feel as though the event is happening all over again—you can see, hear, feel and smell the same things. It is also an unprocessed memory. See page 33 for more detail about flashbacks.

The memory

Memories of sexual violence can be distressing. Sexual violence is a highly traumatic experience. The memories can be:

- **Intrusive**
The memory keeps appearing because it is unfinished business, and you really wish you could somehow finish the experience
- **Fragmented**
Bits and pieces are remembered

- **Sensory**
The pictures, sounds, smells and feelings during a flashback are so clear that the event is remembered by all of, or a combination of some senses
- **Non-sequential**
You can't remember which event happened first
- **Non-verbal**
It is difficult to put into words
- **Non-contextual**
The main aspects are remembered, but it's harder to remember where / when etc
- **Buried**
It may not be remembered until years later, it may be revealed in a dreamlike quality, in snapshots, or ultra-vivid such as flashbacks.



Some ways that you can deal with your thoughts

Keeping a journal can be invaluable for survivors. It can be a good way of managing and exploring your thoughts. It can be the first step to you regaining control. Here are some helpful ways to make the most of your journal.

Bring your thoughts outside.

Writing thoughts down shifts them from the inside to the outside. Now you can explore them and think about whether they are helpful.

Consider triggers.

(Keep yourself safe while doing this). Can you figure out what, when, where and who can set off a flashback or anxious thought? A trigger record will help you to manage or avoid trigger-situations. It will also help you to prepare for them.

Think about what's important to you.

Make a list of valuables that keep you going on the journey of life. It might include:

- **Inner resources** (e.g. good memories and experiences, previous successes, positive characteristics and strengths, spiritual/ religious beliefs)

- **Outer resources** (e.g. people who love and believe in you, nature, animals, objects, a safe place, activities, aspirations)
- **List your resources.**

Invent your own 'first aid kit'.

Make a list of everything that makes you feel good or helps you relax. Maybe having a long bath, going for a walk in the country or talking to friends. Use this list for inspiration when you're having a bad day.

Draw an image or make a collage.

That can help you when you feel gloomy or sad. Put the poster up where it can inspire you.

>>

Letters of my name.

Write your name in big letters. Describe yourself by using the letters of your name. Every letter must represent something positive about you (abilities, likes, etc.).

Example; Sympathetic, Affectionate, Musical.

Positive things diary.

Note down three positive things that have happened each day before you go to bed.

1. I visited my favourite place,
2. I finally did that pile of washing up,
3. I got through a day without a flashback,
4. I managed to get up and dressed.

Find or make your mantra.

A mantra is a personal power statement of hope and possibility. Tell it to yourself daily; use it to inspire or encourage yourself or to quiet positive thoughts.

Here are some examples:

- “I can do this.”
- “I am kind to myself.”

- “Grant me the serenity to accept the things I cannot change, the courage to change the things I can and the wisdom to know the difference.”
- “This too shall pass.”

My best person.

Is there a person that you know (or used to know) who always supported and believed you? Keep a photograph or any symbol of this person with you always. Refer to it (touch, look at) when you need them.

My best person-response.

Think about what your best person would say to you if you had an unhelpful thought or experience.

What would I say?

If someone close to you said the same thing as you were thinking, what would you say to support them?

Distractions.

Make a list of distractions.

- Contact a friend
- Watch a favourite film or TV programme
- Do something creative
- Dance
- Make yourself a safe cocoon in bed with your duvet

>>

- Visit a place that makes you feel safe (such as a friend or family member's house, a park or beauty spot, a museum/gallery, a cafe)
- Cuddle or play with a pet
- Do some volunteering (some find it comforting to help people, animals or the environment during their own healing process).

Give yourself a calming sleep routine.

Keeping a nightlight on and listening to the radio or calming music may help.

- Electronic screens do not help with relaxation.
- Avoid alcohol, it will stimulate you instead of relaxing you due to its sugar levels. Alcohol is also a depressant which can be unhelpful the following day.
- Slow your breathing, practice the 7:11 breathing technique.
 - Count in for 7 and out for 11. It may take a few minutes but the body will respond regardless of what your mind is thinking.

Don't blame yourself.

If you find yourself thinking 'Why didn't I...', 'It was my fault', 'I should have...', ask yourself:

- What is the evidence?
- If someone told me the same story, who would I blame?
- Are there other ways to think about what happened?
- What are the positives and negatives of this way of thinking?
- What logical errors am I making?
- What would my 'best person' say to tell me that it wasn't my fault?
- What would an organisation like RSACC say?

You have survived and you are healing. Be kind to yourself, you deserve it

If a friend or a family member told you that this had happened to them would you blame them? Would you think they should have reacted differently? Would you think of them any differently?

Feelings

Your emotional reactions

When dealing with trauma, you can feel either a huge mass of feelings that are difficult to separate, or an empty numbness.

Survivors sometimes feel overwhelmed by many different feelings. They might feel them in their throat, or their stomach, or somewhere else. They may be experienced as a mass of feelings that are hard to describe.

When you look at this huge block of feelings, you will notice some individual ones. Often it is helpful to explore these feelings, identify each one, and investigate the thought behind it.



Begin to look at your feelings individually and explore them:

Anger is not the enemy.

“Bitterness is like cancer. It eats upon the host. But anger is like fire. It burns it all clean.”

(Maya Angelou)

Anger can show you that what is important to you is under threat. Anger protects you from future betrayal. It warns you that you must do something about the reason behind it. The way a person chooses to deal with anger can sometimes be a problem, but not the feeling itself.

Shame is one of the most destructive emotions.

“Shame corrodes the very part of us that believes we are capable of change.”

(Brené Brown, *I Thought It Was Just Me: Women Reclaiming Power and Courage in a Culture of Shame*)

People usually feel shame because they feel they are not good enough or that something is wrong with them. Sadly, shame can make survivors fear talking to people who could be supportive and who could help them. Survivors do not deserve to feel ashamed.

Loneliness.

“The worst part of holding the memories is not the pain. It's the loneliness of it. Memories need to be shared.”

(Lois Lowry, *The Giver*)

Many survivors feel they are unable to trust anyone. They might be afraid of sharing their experiences with others. They might feel as though they are alone that no-one understands how they feel or what they went through.

Emptiness.

“In all our searching, the only thing we've found that makes the emptiness bearable is each other.”

(Carl Sagan)

The feelings are sometimes so overwhelming that in order to cope a survivor buries them. These are called dissociated emotions. You may worry that other people will blame you, or you may have been told it was your fault. This is absolutely not true. Sexual violence is always the fault of the abuser.

There are people who understand these feelings and want to support you.

Some ways of dealing with emotional reactions

Set yourself small and manageable targets:

- Manage the levels of your feelings by grounding them with an object (see page 20)
- Express your feelings in a safe environment
- Identify and name each one of your feelings
- Try to let them out in the open, write or talk about them
- Accept that your feelings are allowed, and are normal. Each of them has a cause. Each of them belongs to you
- Sort them as belonging to either the past or to the present
- Look at the thought that is behind each feeling.

Listen in and out.

Close your eyes and identify the furthest sound you can hear. After 30 seconds bring your listening in and identify the closest sound you can hear. This may include sounds from your own body like your breathing or your heartbeat. Repeat.

Take your feelings outside yourself.

You can draw a map of your feelings with a colour representing each feeling. Think about the size of your feeling, and give it a similar amount of room on the paper. Think about the thoughts behind the biggest feelings and where they've come from.

Use alone time to learn or practise a skill or hobby.

If you don't feel ready or aren't always in the right mood to be around others, filling your time with hobbies can help ease feelings of loneliness. For example, learning a language, a craft or a musical skill are hobbies that can be practised alone. YouTube videos can be good for this.

Try counselling or face-to-face support.

This can be a good way of helping to express and process feelings such as anxiety, fear, anger, depression, guilt, shame and low self-worth. Learning to share your feelings and experiences with a trained professional or non-judgmental trusted person can help you to feel understood and supported, and to accept and value yourself.



Meditation and Relaxation

“The thing about meditation is you become more and more you.”

David Lynch

Managing yourself on a daily basis can be an overwhelming task and it’s only natural that at times escaping from your own experience is needed.

Meditation, mindfulness and relaxation all have one thing in common. They help your mind, body and soul relax, take a breather and escape for a while. By doing this you are resting body and mind, giving yourself a chance to stop.

Simply taking 5 minutes a day to stop can be very helpful. It takes practice and your mind will wander but that’s ok. Keep trying it.



Here is an example of a very basic meditation.


- Breathe deeply in and out
- Seek to be aware of the different physical sensations you're experiencing as you breathe deeply in and out
- Notice how your body expands with an inhalation and contracts with exhalation
- Experience the breath coming in through your nose, out through your mouth and experience how it feels through other parts of your body
- Notice the stillness around you before and after each breath
- Come back to your breath whenever the thoughts try to take over.

You could consider breathing in a colour. Maybe a clean calm colour such as a **light blue.** With each additional breath in, imagine the colour spread through all your body. Beginning with your head and chest, through your arms into your fingers and down into your legs and your toes.

Then consider the out breath. On each outbreath imagine your breathing out another colour. Maybe a brown or murkier colour representing all the stress and pressure leaving your body. Imagine that colour being breathed out from your toes, up your legs, through your chest, from your arms.

Further examples can be found on many internet sites, such as YouTube. There are also mobile phone Apps that can be useful (e.g. Headspace, Smiling Mind).

www.wikihow.com/Meditate-for-Beginners



Grounding step-by-step

Guidelines

1. Grounding can be done anywhere, any place or any time and no one needs to know you are doing it.
2. Use grounding when you are experiencing a trigger, when you feel strong emotions, feel like using substances, harming yourself or feel yourself dissociating.
3. You can rate your mood before and after, on a 0-10 scale.
4. Keep your eyes open to stay in touch with the present.
5. Stop yourself from talking about negative feelings at this stage, you want to distract away from this.
6. Focus on the here and now, not the past or future.
7. Grounding is more than just a relaxation strategy, it is used to distract and help extreme negative feelings. It is believed to be more effective for trauma than relaxation alone.

How to do it

Mental Grounding

1. **Have a good look around and describe your environment in detail**, e.g. *'I am on the train, I can see trees and a river....'*
2. **Mental games**, e.g. go through the alphabet thinking of different things such as types of dogs, cities etc.
3. **Age progression**, if you have regressed you can slowly go back up e.g. I am now 9, 10 etc. until you are back up to your current age.
4. **Describe an everyday activity in detail**, such as how to follow a recipe.
5. **Imagery**, for example imagining a stop sign in your head, gliding on skates away from the pain, changing the 'TV channel' in your head to a better 'show' or imagining a wall as a buffer between you and the pain.
6. **Safety statements**, thinking *'I am safe now, I am in the present not the past, I am in this location and the date is...'*
7. **Use humour**, think something funny.
8. **Use concentration**, say the alphabet backwards or practise a tricky sum.

Physical Grounding

1. **Run warm or cool water over your hands.**
2. **Focus on your breathing,** notice each inhale and exhale, slow it down and repeat the word safe on each inhale.
3. **Grab tightly onto your chair as hard as you can.**
4. **Touch different objects,** your pen, your keys etc.
5. **Dig your heels into the floor;** remind yourself that you are connected to the ground.
6. **Carry a grounding object in your pocket,** a small rock etc. in your pocket that you can touch whenever you feel triggered.
7. **Stretching,** extend your arms, fingers or legs as far as you can.
8. **Clench and release your fists.**

Soothing Grounding

1. **Say kind statements to yourself,** e.g. you will get through this etc.
2. **Picture people you care about,** look at photos of them.
3. **Think of a safe place,** it could be real or imagined, for example the beach, mountains etc.
4. **Say coping statements** such as I can handle this, I have done it before etc.
5. **Plan a safe treat** such as a nice dinner, relaxing bath etc.
6. **Think of things you are looking forward to,** like seeing a close friend.



What if grounding doesn't work

People who have used grounding say it does work but requires practise to make it as effective as possible. The more you practise it the better it will work, so try to do some every day, it will become automatic after a while.

You don't have to use the methods listed on the previous pages, you could think up your own method, you may find that it works better for you. Try to start grounding as early as possible in a negative mood cycle, for example just after a flashback, don't leave it until later.

You could create a recording of a grounding message that you can play whenever you need it, if you don't want to use your own voice you could ask someone close to you to help.

You can also teach family and friends about grounding so they can help if you become overwhelmed.

Notice which method works best for you, you can use the mood rating scale for this, and lastly, don't give up!



Body

Your body's reactions

Our bodies can hold the trauma from rape or sexual abuse. Every survivor responds differently to what happened to them.

- 1. Look at your body's unique responses, and any responses that others have also had**
- 2. Recognise which bodily reactions are related to the abuse, and why your body decided to react in that way**
- 3. Think/talk about if these reactions are helpful or not right now. Decide whether you are going to deal with the bodily reaction(s).**



Some common bodily reactions to rape or abuse are:

- Irritable Bowel Syndrome
- Pains
- Tensions
- Headaches
- Changes in body language

Becoming aware of changes and patterns in any bodily reactions are a good first step towards controlling them.

Don't expect too much of yourself at once. There are many different ways to work on your body – anything from yoga to massage to strengthening exercises. If you feel something isn't working for you, review it and try a different approach.



Some ways to look after your body

A stronger body can help you deal with your system being overactive.

Where in your body do you feel the most pain?

Is there anything you can do to soothe it, either with the aid of medicine, advice from a healthcare professional (for example your GP, NHS Direct, a walk-in clinic), or things you can do at home?

If you are not comfortable seeing a male doctor you have the right to request a female doctor. Also you can refuse any examinations you are uncomfortable with. All professionals should ask your permission to examine you or touch you.

Some ways you can soothe your body:

Release positive chemicals through exercise.

Frequent and regular physical exercise boosts the immune system, improves mental health, helps prevent depression, and encourages positive self-esteem. Exercise is also good for sleep.

Strength training.

Being stronger helps you feel more in control of your body and environment.



Give yourself time to mend:

Sleep.

Turning your computer, TV or tablet off 45 minutes before sleeping helps to relax the mind and prepare it for sleep.

Laughter.

Watch a favourite comedy. Laughter releases endorphins, which are our body's natural painkillers. When you laugh your whole body relaxes.

Think about how you could cut down on alcohol, caffeine, sugar or cigarettes.

These things can make you feel good for a short time and finding things that make you feel good is healthy and normal. Identifying things that might make us feel worse in the long run and reducing them can be good for your body and mind.

Relaxation techniques may not always help with post-traumatic stress disorder. If trying to relax is making you more anxious; you could try exercise that helps you release energy and build strength.

Behaviour

Your actions

When someone has suffered any kind of rape or abuse it can affect their behaviour. There are very good reasons for a survivor to behave in certain ways. There will always be some thoughts or feelings behind our behaviour:

Self-protective thoughts or feelings behind your behaviour:

- Fear that this secret, which is often or always on their mind, might slip out
- Thoughts that others may reject them
- Fear of being questioned
- A feeling that they can trust nobody
- Trying to control anxiety or exposure to triggers
- Trying to block out memories
- Trying to 'keep the box shut' by distracting themselves
- Fear of seeing the abuser somewhere
- Fear of having an anxiety attack in front of others.



In order to change your behaviours, you will need to believe it is important enough to make that change, and have confidence in your ability to make the change.

Identifying the thoughts and feelings behind the behaviour will help you to understand and work on your reactions as a whole. Some of these thoughts and feelings can help survivors to cope and to process what has happened safely. Others may cause the survivor further distress or be harmful.

- Write down some of the behaviours that you find yourself doing
- Ask yourself what the reasons behind those behaviours are
- Think about how they, and the behaviours they are causing, are making you feel in the short and long-term
- Evaluate them; are they helpful or unhelpful?

Pushing people away

...When I push people away I feel more in control but more alone.

Pushing people away can be an instinctive way to protect yourself from further harm. Isolation can often reflect how a survivor acted during or after rape or abuse. The perpetrator might have forced you to keep it a secret by making threats or telling you it's normal or that nobody will believe you. It's normal to be cautious to trust people again. Finding other people to talk to can help you to think about your own thoughts and reactions. Others can tell you when your thoughts are unhelpful or self-destructive, and can help you to see that what happened wasn't your fault.

Not wanting to be intimate

...How will I ever trust anyone again? I feel like my body isn't my own.

A lot of people fear intimacy after rape or abuse. Take your time, do things as you are ready. What happened to you was not sex. Sex is different because you are in control. It's important your partner understands and allows you to take it at your own pace.

Promiscuity

...I want to have control back over my body. Sometimes, I don't feel how I expected to afterwards.

Some people find that they want to have lots of sex, sometimes with lots of different people. This can be a way to take back control of your body. Sex is a healthy and natural thing, but be aware if you are putting yourself in some risky situations. Make sure you are safe and think about how it makes you feel.

Being very vigilant

...I am constantly on edge.

Feeling over-attentive and alert to every sound and movement is normal after trauma because in that way you protect yourself against danger. As you work through the feelings and thoughts behind this vigilance, it will become easier.

In the short-term, try to find a safe place that is able to soothe you and go there once a day.

Not wanting to go out

...I don't want to see him, or anyone else. I don't have the energy to go out anyway.

Your home can be a therapeutic place where you may feel you're safest. In the short-term, it might enable you to get through the day. Try to think about what you are compromising in the long-term. Are there any other ways of achieving these things in small, manageable and safe ways?

Drinking / smoking more

...It makes me calmer, helps me to be sociable.

These are normal coping mechanisms in any stressful situation. In what situations are you drinking or smoking? How do you feel before and afterwards? Is it leading to any risky or challenging situations?

Being more irritable

...I keep snapping at people. It's not their fault and I'm pushing them away.

Everybody is irritable sometimes. Extreme irritability can come as a result of a lack of sleep, feelings of isolation or lack of control amongst other things. At what times of the day are you irritable?

“If we can share our story with someone who responds with empathy and understanding, shame can't survive.”

Brené Brown, *Daring Greatly: How the Courage to Be Vulnerable Transforms the Way We Live, Love, Parent, and Lead.*

Reaching out to others, however difficult, can be really helpful

Isolation and secrecy often reflect how a survivor acted during or after rape or abuse. You might have never told anybody for fear of what they might think or say or fear of not being believed. Being with others can distract you and comfort you.

Finding other people to talk to can help you to think about your own thoughts and reactions. Others can tell you when your thoughts are unhelpful or self-destructive, and can help you to see that what happened wasn't your fault.

As well as identifying the behaviours that are associated with recovery, and thoughts and feelings behind them, creating new positive and therapeutic behaviours can be helpful. Here are some ideas:

- Call, write to, email or visit a friend or member of the family
- Give a genuine compliment
- Think about what your friends and family give or mean to you – and tell them
- Join a group (singing, sport, craft, wellbeing)
- Do something nice for a stranger
- Donate to a worthy charity
- Create a /digital playlist, artwork or photo album for a friend
- Print out an inspirational poem
- Be kind to yourself

“If we change within, our outer life will change also.”

Jean Shinoda Bolen



It's never too late to talk

Understanding your reactions to the abnormal situation you have been in is the first step towards starting to take back control and finding someone to talk to. Many survivors do not feel they can talk to their family or friends about what happened or how it has affected their thoughts, but this does not mean you are alone.


Having a supportive network around you, finding someone to talk to who you trust and will listen - **will help the healing.**



Other support networks.

Where else could you go for help if you wanted to? Other ways to get support could be:

- Talking to a trusted friend or family member. Choose this person carefully: do you trust them enough, will they show you kindness and try to understand you?
- Online survivors; groups and forums
- Sexual violence survivors' meetings/groups
- Learn a new skill in a new environment
- Counselling, through the NHS via your GP, or via a specialist service
- Medication such as anti-depressants - Consult your GP



Who could you talk to?

Key reactions: Flashbacks

It is common for a survivor to experience vivid images of the rape or sexual abuse, making you feel as though the event is happening all over again. These vivid images are known as flashbacks and are active memories of a past traumatic event. Flashbacks can occur at any time, are out of a person's control and difficult to get rid of.

Though flashbacks are horrible and terrifying, they can show that a person is ready to deal with the traumatic event.



Some ways you can help yourself to cope with flashbacks:

- 1. Tell yourself that you are having a flashback:**
Learn to recognise what happens to your body when you are having a flashback. Recognise the clues your body gives you just before a flashback happens.
- 2. Remind yourself that the worst is over:**
The feelings and sensations you are experiencing now are memories of the past. The actual event took place in the past and you survived it.
- 3. Get grounded and focus on the present:**
Use all your senses to keep you focused on the here and now. Look around and notice different colours, objects around you. Listen to the different sounds - music, people or your own breathing. Feel your body, your clothes, the chair or the floor supporting you. Stamp your feet on the ground, so that you know where you are and that you are no longer trapped in a situation you cannot escape from.

- 4. Breathe:**
When you are scared it is likely that you may experience shallow breathing or stop breathing altogether. As a result your body could react to the lack of oxygen, causing panicky feelings, a pounding heart, pounding in the head, tightness, sweating, faintness and dizziness. If you breathe deeply some of this panic will stop. Put your hand on your stomach and breathe in so that your stomach pushes against your hand and then breathe out so that the stomach goes in.

- 5. Compare then and now:**
Remind yourself of where you were then and where you are now. *'I was in the sitting room then. Now I am in my bed.'*

>>



6. Give yourself some boundaries:

Sometimes when you are having a flashback you might lose the sense of where you finish and where the world begins: as if you do not have skin. Wrap yourself in a blanket, hold a pillow or go to bed; do anything to help you to feel protected.

7. Tell your friends, partner or relatives about flashbacks:

It's important that the people around you know about flashbacks so that they know how to help you. You may want someone to be with you during or just after a flashback, or you may prefer to be alone. Decide what you find most helpful and let others know.

8. Take time to recover:

Flashbacks are very powerful, so give yourself some time afterwards. Don't expect yourself to jump into activities right away. Take a nap, have a warm bath or take some quiet time. Show yourself kindness, gentleness and patience, allow your body some comfort.

9. Bring your memories into the open:

Anyone who understands the effects of rape, sexual assault and childhood sexual abuse can be a valuable resource to helping you heal. If you feel able, it can be helpful to write down the sights, sounds, smells, emotions and sensations from your flashback too. Putting it on paper can help to get it out of your mind.

10. Finally, know you are not crazy.

You are healing and you're dealing with the normal reactions to a traumatic event.

These facts and suggestions have been adapted from:
Rape & Sexual Violence Project
www.rsvporg.co.uk

Key reactions: Self-harm

What is Self-harm?

Self-harm is a broad term that describes the act of hurting yourself on purpose. It is also known as self-inflicted violence, self-injury or self-mutilation.

Self-harm could include any of the following behaviours:

- Cutting, burning or slashing the skin
- Obsessive cleaning of the skin, sometimes using bleach
- Pulling hair out
- Repeatedly hitting the body

People can also harm themselves by drinking alcohol, smoking or taking drugs (all of which could lead to addiction), eating too much/too little, (which could lead to eating disorders), like anorexia, bulimia or compulsive eating or work extremely hard, leading to exhaustion).



Who carries out self-harm?

Self-harm is carried out by women and men; there is no typical person who self-harms and people of all ages, backgrounds and cultures can self-harm.

Why do people self-harm?

Self-harm can be a way to:

- Express, cope with or numb painful feelings
- Get some control over painful and often confusing feelings
- Punish yourself; this can be particularly true if a person has low self-esteem or blames themselves for the original trauma
- Provide comfort at a difficult time by doing something that is familiar. This can help when new and confusing feelings appear

Many people who self-harm feel hopeless or worthless at the time. As a result they may ignore and/or deny their own needs. If their emotions seem impossible to understand or deal with then destroying those feelings might seem like the only answer.

Myths about self-harm:

“It is a failed suicide attempt”

Self-harm is much more about attempting to cope with life rather than wanting to die. Injuries can be life threatening but are rarely so.

“Self-harm is attention seeking behaviour”

Many people try as hard as they can to hide any evidence of their self-harm. People do not hurt themselves to gain attention.

“Self-harm is a sign of someone going mad”

Self-harm is a sign of deep distress – not madness.

“People who self-harm are a danger to others”

People who self-harm are directing the hurt at themselves and not at other people. In fact, most people who self-harm would be appalled at the idea of harming anyone else.



What can help?

Beginning to understand the reasons for and the meaning of your self-harm will be useful. Keep a diary and ask yourself the following questions:

How do you feel before and after the self-harm?

Asking yourself this question might help you understand the feelings you are trying to cope with. It may be easier to find another way to deal with those feelings once you have identified them.

What does self-harm mean to you?

What do the feelings signify, in your past or your life at the moment? When did you first start to self-harm? Identifying these meanings can help you to understand what it gives you.

Why do you want to stop hurting yourself?

Is it due to pressure from others, a desire to reduce your feelings of shame and guilt about the self-harm or have you decided that self-harm doesn't work for you anymore?

What resources do you have to support you if you stop or reduce your self-harming?

Have you got people around you who know about the self-harm and want to support you? Think about what you need from these people and how they help you.

When do you normally self-harm? Is there a pattern?

For example, you always self-harm in the evening or you always self-harm when you are left alone in the house. Changing a ritual or thinking ahead about certain situations might have an effect on the self-harm in general.

Replacement and distraction are useful tools. Can you make a list of things that you can do instead of self-harm? For example, call a friend or do something physical (e.g. exercise) to release the feelings you have inside.

Breaking it down

On these pages you can note down your favourite self-help suggestions from each section.

Thoughts

Feelings

Body

Behaviour

Which parts of the self help guide have you found particularly useful?

Do you have any other ideas that aren't mentioned?

We'd really like to know your opinions and tips so that we can share

Further resources

There are many resources for you to use. These are just a few we have found.

**Beginning To Heal:
A First book for Men & Women
who were Sexually Abused
as Children**
By Ellen Bass and Laura Davies

**Quest for Respect:
Healing Guide for Survivors
of Rape**
By Linda Braswell

**New Shoes:
Stepping Out of the Shadow
of Sexual Abuse & Living Your
Dreams**
By Rebecca Mitchell

**The Sexual Healing Journey:
A Guide for Survivors of
Sexual Abuse**
By Wendy Maltz

**The Courage to Heal:
A Guide for Women Survivors
of Child Sexual Abuse**
By Ellen Bass and Laura Davies

The Courage To Be Me
By Nina Burrowes
Book available online:
[https://ninaburrowes.com/
books/the-courage-to-be-
me/preface](https://ninaburrowes.com/books/the-courage-to-be-me/preface)

Panic Attacks
www.nomorepanic.co.uk
[http://www.rapecrisisscotland.
org.uk/publications/RCS_
supportresources_panicattacks.
pdf](http://www.rapecrisisscotland.org.uk/publications/RCS_supportresources_panicattacks.pdf)

Pandora's Project
Online support and
resources for survivors.
www.pandys.org

Eating Disorders
www.beateatingdisorders.org.uk
[www.nhs.uk/conditions/
eating-disorders.](http://www.nhs.uk/conditions/eating-disorders)
www.niwe.org.uk

Meditation and Relaxation

Online self-help site.
www.headspace.com

Tracks on YouTube
e.g. Meditation relax club

Phone Apps: Headspace,
Smiling Minds, Meditation
Relaxation Club.

PODS

For information on dissociation:
www.pods-online.org.uk
[www.firstpersonplural.org.uk/
about-first-person-plural](http://www.firstpersonplural.org.uk/about-first-person-plural)

Nightmares and Sleep Problems

For information and self-help
[http://www.rapecrisisscotland.
org.uk/publications/RCS_supp
ortresources_nightmares.pdf](http://www.rapecrisisscotland.org.uk/publications/RCS_supportresources_nightmares.pdf)

Rape, Abuse and Incest National Network

Online network and
information for survivors.
www.rainn.org

Self-Harm

Online resources for self-harm
support and self-help.

<https://www.selfharm.co.uk>

[http://www.mind.org.uk/inform
ation-support/types-of-mental-
health-problems/self-
harm/#.VRABefmsVcc](http://www.mind.org.uk/information-support/types-of-mental-health-problems/self-harm/#.VRABefmsVcc)

RSACC

Contact Details:

**The Rape & Sexual Abuse
Counselling Centre
(Darlington & County
Durham) CIO**

Charity Address:
PO Box 106
Darlington DL3 7YS

Charity No: 1164121

Business Line:
01325 354 119

Helpline:
0300 222 5730
(check social media
for days and times)

Lines are open:
Monday to Thursday
from 6.30pm – 9pm

Email:
info@rsacc-thecentre.org.uk

Website:
www.rsacc-thecentre.org.uk



Contact us

Phone: 01325 354119

Email: info@rsacc-thecentre.org.uk

www.rsacc-thecentre.org.uk

PO Box 106, Darlington,
Co. Durham DL3 7YS

Charity No. 1164121



RSACC ♀
Rape and Sexual Abuse Counselling Centre
Darlington and County Durham